

Western Australian Immunisation Schedule

Childhood

Age	Antigen	Vaccine brand(s)	Important information
Birth to <7 days	Hepatitis B	H-B-Vax [®] II Paed or Engerix [®] B Paed	
6-8 Weeks	DTPa-hepB-IPV-Hib	Infanrix [®] hexa	
	13vPCV	Prevenar [®] 13	
	Rotavirus	Rotarix [®]	Oral dose 1: 6–14 weeks of age
Aboriginal	MenACWY	Nimenrix [®]	
Aboriginal and/or MRC	MenB	Bexsero [®]	Prophylactic paracetamol recommended prior to dose
4 Months	DTPa-hepB-IPV-Hib	Infanrix [®] hexa	
	13vPCV	Prevenar [®] 13	
	Rotavirus	Rotarix [®]	Oral dose 2: 10–24 weeks of age
Aboriginal	MenACWY	Nimenrix [®]	
Aboriginal and/or MRC	MenB	Bexsero [®]	Prophylactic paracetamol recommended prior to dose
6 Months	DTPa-hepB-IPV-Hib	Infanrix [®] hexa	
Aboriginal and/or MRC	13vPCV	Prevenar [®] 13	
MRC	MenB	Bexsero [®]	Prophylactic paracetamol recommended prior to dose
12 Months	MMR	MMRII [®] or Priorix [®]	
	13vPCV	Prevenar [®] 13	
	MenACWY	Nimenrix [®] *	
Aboriginal and/or MRC	MenB	Bexsero [®]	Prophylactic paracetamol recommended prior to dose
MRC	Hepatitis B	H-B-Vax [®] II Paed or Engerix [®] B Paed	<32 weeks gestation or <2000g birth weight
All children > 12 months	Children diagnosed with MRC at >12 months of age require 1 dose of 13vPCV at diagnosis, followed by a first dose of 23vPPV, 2 to 12 months later or at 4 years of age, whichever is later. A second dose of 23vPPV is given 5 years later. See Australian Immunisation Handbook		
MRC			
18 Months	DTPa	Infanrix [®] or Tripacel [®]	
	Hib	ActHIB [®]	
	MMRV	Priorix-Tetra [®] or ProQuad [®]	Do not administer MMRV as dose 1 of MMR containing vaccine if <4 years of age
Aboriginal	Hepatitis A	Vaqta [®] Paed/Adol	First dose of the 2 dose schedule
<2 Years	Catch-up for MenB is available for Aboriginal children <2 years of age until 30 June 2023.		
Aboriginal			
4 Years	DTPa-IPV	Quadracel [®] or Infanrix [®] IPV	
Aboriginal	Hepatitis A	Vaqta [®] Paed/Adol	Not required if previously received 2 doses (first dose at age ≥12 months) at least 6 months apart
	23vPPV	Pneumovax23 [®]	First dose (if not previously received) Second dose at least 5 years later

Age	Antigen	Vaccine brand(s)	Important information
All adolescents and adults with MRC	<p>Adolescents and adults diagnosed with MRC require 1 dose of 13vPCV at diagnosis, followed by a first dose of 23vPPV, 2 to 12 months later. A second dose is given at least 5 years later. People of all ages with specified MRC are eligible for MenACWY, MenB and Hib vaccines. See the ATAGI clinical advice for pneumococcal and meningococcal diseases as well as the Australian Immunisation Handbook for more information.</p>		
Year 7	HPV	Gardasil®9	2 doses of Gardasil®9, 6 months apart
	dTpa	Boostrix®	
Year 10	MenACWY	Nimenrix®	Catch-up for adolescents <20 years of age
Pregnant	dTpa	Boostrix® or Adacel®	Each pregnancy, ideally between 20–32 weeks
≥20 Years Aboriginal	Hepatitis B	Engerix-B (adult)	For non-immune persons. See the Australian Immunisation Handbook for recommendations and timing of doses
≥50 Years Aboriginal	13vPCV 23vPPV	Prevenar® 13 Pneumovax23®	See the Australian Immunisation Handbook for for timing of doses
Born after 1965	MMR	MMRII® or Priorix®	Give 1 or 2 doses as required
≥70 Years	13vPCV	Prevenar® 13	
≥70 to 79 Years	Herpes-zoster	Zostavax®	Catch-up program for 71–79 year olds until 31 Oct 2021

2021 Seasonal influenza

Age	Vaccine brand(s)	Important information
6 Months to <5 Years	Vaxigrip® Tetra, Fluarix® Tetra	If child <9 years, 2 doses required 1 month apart in the first year they receive influenza vaccine
≥5 Years to Year 6	FluQuadri® (WA State Program)	
5 to <65 Years Aboriginal and/or MRC	Afluria® Quad, Fluarix® Tetra, Vaxigrip® Tetra	
Pregnant	Afluria® Quad, Fluarix® Tetra, Vaxigrip® Tetra	Each pregnancy, any trimester
≥65 Years	Fluad® Quad	Use standard QIV if Fluad® Quad unavailable
WA Health staff	Fluarix® Tetra, FluQuadri® (WA State Program)	For ≥65 years use Fluad® Quad

The term 'Aboriginal' is inclusive of all Aboriginal and Torres Strait Islander people.

MRC refers to medical risk conditions. Refer to the current online edition of The [Australian Immunisation Handbook](#) for all medical risk conditions.

Persons aged <20 years can catch-up scheduled vaccines, refer to [Australian Immunisation Handbook](#) for timing of doses.

Adult refugees and humanitarian entrants are eligible for dTpa, Hepatitis B, IPV, MMR and VZV.

* **MenACWY catch-up** for Aboriginal children <5 years of age until 31st January 2023.