### **VAX** vaccinate your baby

- Needles are safe and the best way to protect your baby
- lust because your baby has had their needles doesn't mean they are fully protected. Needles only cover the most common strains. That's why it is important to know the symptoms!
- The needles listed in blue are extra ones that are only given to Indigenous babies like yours, who need them because they are more at risk

#### Vaccination Schedule for Meningitis, **Meningococcal and Pneumococcal**

Age	Type of Vaccination
2 Months	<ul> <li>Haemophilus influenza type B (Hib)</li> <li>Pneumococcal</li> <li>Meningococcal B - only Aboriginal and Torres Strait Islander (ATSI)</li> </ul>
4 Months	<ul><li>Haemophilus influenza type B (Hib)</li><li>Pneumococcal</li><li>Meningococcal B - only ATSI children</li></ul>
6 Months	<ul><li>Haemophilus influenza type B (Hib)</li><li>Pneumococcal - only ATSI children</li></ul>
12 Months	<ul><li>Meningococcal ACWY</li><li>Pneumococcal</li><li>Meningococcal B - only ATSI children</li></ul>
18 Months	<ul> <li>Haemophilus influenza type B (Hib)</li> </ul>

"You would rather get the injection now than have them in hospital waking up with limbs missing from the disease itself"

- Mrs Wallace from the APY region

"Vaccinating Aboriginal and Torres Strait Islander children is of great importance" - Dr Tanya Schramm, a Palawa woman

> Every Second Counts!

For further information visit: www.meningitis.com.au



**MENINGITIS CENTRE AUSTRALIA** 

Free Call 24/7 1800 250 223



info@meningitis.org.au



artwork designed by Tyrown Waigana. 2020 NAIDOC poster designer



Meningitis, Meningococcal and Pneumococcal Keeping Bubba Safe

**ACTVAX** 

.....







# Hi. This is important information for your mob

Meningitis, Meningococcal and Pneumococcal are VERY SERIOUS illnesses, especially for babies.





1 in 5 survivors will have disabilities like brain damage, loss of arms or legs, or deafness

"We've seen so many cases of meningococcal in town, I wanted to make sure my kids wouldn't get it too" – Sho-Ella Smith



## KNOW the Symptoms - Danger!



High Temperature



**Throwing Up** 



Sleepy



Blotchy Skin/Rash



Blank Eyes / Doesn't like being Touched



Light Hurts Eyes



Cold Hands and Feet



Sore Neck and Back Arching



**Not Hungry** 



Hard to Breathe



Loud Cry, Bump on Top of Head

- Meningitis can kill within 24 hours. The time between the first symptom and your baby being very sick may only be a few hours
- All babies will show different signs, so you need to know them all- check your fridge magnet!

#### **KEEP YOUR MOB SAFE**

## **ACT** immediately

- If your bubba has any of these symptoms, go to a doctor or hospital straight away
- Ask the doctor- 'could it be Meningitis, Meningococcal or Pneumococcal?'
- Don't wait for a rash to appear on bubba.
   It's the symptom most people know but some babies might not get it at all
- Trust your gut as a parent. You know your bubba better than anyone else. If you get sent home from the hospital but baby gets worse, go straight back

